

数式処理実習pair試験予行演習問題、西谷@関西学院大・
理工、
2016/6/3 実施

▼ 1(1)

```
> restart;
simplify(diff(sqrt(x^2+1)*(x^3+1)^(1/3),x));
```

$$\frac{x(2x^3 + x + 1)}{\sqrt{x^2 + 1} (x^3 + 1)^{2/3}} \quad (2.1)$$

▼ 1(2)

```
> restart;
x:=t->a*cos(t)^3;
y:=t->a*sin(t)^3;
```

$$x := t \rightarrow a \cos(t)^3$$

$$y := t \rightarrow a \sin(t)^3 \quad (3.1)$$

```
> dxdt:=diff(x(t),t);
dydt:=diff(y(t),t);
```

$$dxdt := -3 a \cos(t)^2 \sin(t)$$

$$dydt := 3 a \sin(t)^2 \cos(t) \quad (3.2)$$

```
> dydt/dxdt;
```

$$-\frac{\sin(t)}{\cos(t)} \quad (3.3)$$

▼ 2(1)

```
> restart;
eq1:=1/(cos(x)^2+4*sin(x)^2);
```

$$eq1 := \frac{1}{\cos(x)^2 + 4 \sin(x)^2} \quad (4.1)$$

```
> diff(eq1,x);
```

$$-\frac{6 \cos(x) \sin(x)}{(\cos(x)^2 + 4 \sin(x)^2)^2} \quad (4.2)$$

```
> int(eq1,x);
```

$$\frac{1}{2} \arctan(2 \tan(x)) \quad (4.3)$$

▼ 2(2)

```
> restart;
eq2:=1/((y-x)^(1/3));
```

$$eq2 := \frac{1}{(y - x)^{1/3}} \quad (5.1)$$

```
=> eq3:=int(int(eq2,x=0..y),y=1/n..1);
```

$$eq3 := -\frac{9}{10} \left(\frac{1}{n}\right)^{5/3} + \frac{9}{10} \quad (5.2)$$

```
=> limit(eq3,n=infinity);
```

$$\frac{9}{10} \quad (5.3)$$

▼ 3(1)

```
> restart;
with(LinearAlgebra);
> A:=Matrix([[1,-1,1],[-7,2,1],[2,1,2]]);
```

$$A := \begin{bmatrix} 1 & -1 & 1 \\ -7 & 2 & 1 \\ 2 & 1 & 2 \end{bmatrix} \quad (6.1)$$

```
=> l,P:=Eigenvalues(A);
l, P := \begin{bmatrix} 3 \\ -2 \\ 4 \end{bmatrix}, \begin{bmatrix} 0 & -1 & -1 \\ 1 & -2 & 4 \\ 1 & 1 & 1 \end{bmatrix} \quad (6.2)
```

```
=> MatrixInverse(P).A.P;
```

$$\begin{bmatrix} 3 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & 4 \end{bmatrix} \quad (6.3)$$

▼ 3(2)

```
> restart;
with(LinearAlgebra);
> A:=Matrix([[0,c,b],[c,0,a],[b,a,0]]);
```

$$A := \begin{bmatrix} 0 & c & b \\ c & 0 & a \\ b & a & 0 \end{bmatrix} \quad (7.1)$$

```
=> B:=Matrix([[0,1,1],[1,0,1],[1,1,0]]);
```

$$(7.2)$$

```
> Determinant(A.B);
```

$$B := \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

(7.2)

$4 \ a \ b \ c$


(7.3)

4(1)

```
> restart;
f1:=x->x;
f2:=x->-3*x+12;
```

$$\begin{aligned} f1 &:= x \rightarrow x \\ f2 &:= x \rightarrow -3x + 12 \end{aligned}$$

```
> plot([f1(x),f2(x)],x=0..5);
```



```
> eq1:=int(f1(t),t=0..x);
g1:=unapply(eq1,x);
```

$$eq1 := \frac{1}{2} x^2$$

$$g1 := x \rightarrow \frac{1}{2} x^2$$

(8.2)

```
> eq2:=int(f2(t),t=3..x)+g1(3);
g2:=unapply(eq2,x);
```

$$eq2 := -\frac{3}{2} x^2 - 18 + 12x$$


$$g2 := x \rightarrow -\frac{3}{2} x^2 - 18 + 12x$$

(8.3)

```
> plot([g1(x),g2(x)],x=0..6);
```

(7.2)

(7.3)



```
> a_l:=unapply(diff(g1(x),x),x);
```

$$a_l := x \rightarrow$$

(8.4)

```
> eq_L:=expand(a_l(a)*(x-a)+g1(a));
```

$$eq_L := -\frac{1}{2} a^2 + ax$$

(8.5)

```
> y_L:=unapply(eq_L,x);
```

$$y_L := x \rightarrow -\frac{1}{2} a^2 + ax$$

(8.6)

```
> Q_x:=solve(y_L(x)=0,x);
```

$$Q_x := \frac{1}{2} a$$

(8.7)

```
> Q:=Vector([Q_x,0]);
```

$$Q := \begin{bmatrix} \frac{1}{2} a \\ 0 \end{bmatrix}$$

(8.8)

```
> eq3:=solve(y_L(x)=g2(x),x);
```

$$eq3 := -a + 6, \frac{1}{3} a + 2$$

(8.9)

```
> R_x:=eq3[1];
```

$$R_x := -a + 6$$

(8.10)

```
> R_y:=expand(g2(R_x));
```

$$R_y := -\frac{3}{2} a^2 + 6a$$

(8.11)

```
> R:=Vector([R_x,R_y]);
```

(8.12)

```

> H:=Vector([R_x,0]);
H :=  $\begin{bmatrix} -a + 6 \\ -\frac{3}{2} a^2 + 6 a \end{bmatrix}$  (8.12)

> eq4:=expand((Q_x-R_x)*R_y/2);
eq4 :=  $-\frac{9}{8} a^3 + 9 a^2 - 18 a$  (8.14)

> S:=unapply(eq4,a);
S := a →  $-\frac{9}{8} a^3 + 9 a^2 - 18 a$  (8.15)

> s2:=solve(diff(S(a),a),a);
s2 :=  $\frac{4}{3}, 4$  (8.16)


> a_max:=s2[1];
a_max :=  $\frac{4}{3}$  (8.17)

```

4(2)

```

> restart;
f1:=x>x;
f2:=x>-3.1*x+12;
f1 := x → x
f2 := x → -3.1 x + 12 (9.1)

> plot([f1(x),f2(x)],x=0..5);

x0:=solve(f1(x)=f2(x),x);
x0 := 2.926829268 (9.2)

> eq1:=int(f1(t),t=0..x);

```

(8.12)

(8.13)

(8.14)

(8.15)

(8.16)

(8.17)

(9.1)

(9.2)

```

g1:=unapply(eq1,x);
eq1 :=  $\frac{1}{2} x^2$  (9.3)


g1 := x →  $\frac{1}{2} x^2$ 

> eq2:=int(f2(t),t=x0..x)+g1(x0);
g2:=unapply(eq2,x);
eq2 :=  $-1.550000000 x^2 - 17.56097561 + 12. x$  (9.4)

g2 := x →  $-1.550000000 x^2 - 17.56097561 + 12. x$ 

```

> plot([g1(x),g2(x)],x=0..6);



> a_l:=unapply(diff(g1(x),x),x);
a_l := x → x (9.5)

```

> eq_L:=expand(a_l(a)*(x-a)+g1(a));
eq_L :=  $-\frac{1}{2} a^2 + a x$  (9.6)

```

```

> y_L:=unapply(eq_L,x);
y_L := x →  $-\frac{1}{2} a^2 + a x$  (9.7)

```

> Q_x:=solve(y_L(x)=0,x);

$Q_x := \frac{1}{2} a$ (9.8)

```

> eq3:=solve(y_L(x)=g2(x),x);
eq3 :=  $-0.3225806452 a + 3.870967742$ 
 $+ 0.000006451612903$  (9.9)

```

$$\sqrt{1.025000000 \cdot 10^{10} a^2 - 6.000000000 \cdot 10^{10} a + 8.780487804 \cdot 10^{10}},$$

$$-0.3225806452 a + 3.870967742$$

$$-0.000006451612903$$

```


$$\sqrt{1.025000000 \cdot 10^{10} a^2 - 6.000000000 \cdot 10^{10} a + 8.780487804 \cdot 10^{10}}$$

> R_x:=eq3[1];
R_x :=  $-0.3225806452 a + 3.870967742$ 
 $+ 0.000006451612903$ 
 $\sqrt{1.025000000 \cdot 10^{10} a^2 - 6.000000000 \cdot 10^{10} a + 8.780487804 \cdot 10^{10}}$ 

```

```

> R_y:=expand(g2(R_x));
R_y :=  $-0.8225806453 a^2 + 3.870967743 a$ 
 $+ 0.000006451612904 a$ 
 $\sqrt{1.025000000 \cdot 10^{10} a^2 - 6.000000000 \cdot 10^{10} a + 8.780487804 \cdot 10^{10}} - 1 \cdot 10^{-8}$ 
 $+ 1 \cdot 10^{-14} \sqrt{1.025000000 \cdot 10^{10} a^2 - 6.000000000 \cdot 10^{10} a + 8.780487804 \cdot 10^{10}}$ 

```

```

> R:=Vector([R_x,R_y]);
R :=  $\begin{bmatrix} -0.3225806452 a + 3.870967742 \\ + 0.000006451612903 \\ \sqrt{1.025000000 \cdot 10^{10} a^2 - 6.000000000 \cdot 10^{10} a + 8.780487804 \cdot 10^{10}} \\ -0.8225806453 a^2 + 3.870967743 a \\ + 0.000006451612904 a \\ \sqrt{1.025000000 \cdot 10^{10} a^2 - 6.000000000 \cdot 10^{10} a + 8.780487804 \cdot 10^{10}} - 1 \cdot 10^{-8} \\ + 1 \cdot 10^{-14} \sqrt{1.025000000 \cdot 10^{10} a^2 - 6.000000000 \cdot 10^{10} a + 8.780487804 \cdot 10^{10}} \end{bmatrix}$ 

```

```

> H:=Vector([R_x,0]);
H :=  $\begin{bmatrix} -0.3225806452 a + 3.870967742 \\ + 0.000006451612903 \\ \sqrt{1.025000000 \cdot 10^{10} a^2 - 6.000000000 \cdot 10^{10} a + 8.780487804 \cdot 10^{10}} \\ [0] \end{bmatrix}$ 

```

```

> eq4:=expand((Q_x-R_x)*R_y/2);
eq4 :=  $-0.5516389180 a^3 + 4.432882415 a^2$ 
 $+ 0.000005306971905 a^2$ 
 $\sqrt{1.025000000 \cdot 10^{10} a^2 - 6.000000000 \cdot 10^{10} a + 8.780487804 \cdot 10^{10}}$ 
 $- 9.319560420 a$ 
 $- 0.00002497398544 a$ 
 $\sqrt{1.025000000 \cdot 10^{10} a^2 - 6.000000000 \cdot 10^{10} a + 8.780487804 \cdot 10^{10}}$ 
 $+ 1.652242329 \cdot 10^{-8}$ 
 $+ 1.290322580 \cdot 10^{-14}$ 

```

(9.10)

(9.11)

(9.12)

(9.13)

(9.14)

$$\sqrt{1.025000000 \cdot 10^{10} a^2 - 6.000000000 \cdot 10^{10} a + 8.780487804 \cdot 10^{10}}$$

```
> S:=unapply(eq4,a);
```

$$S := a \rightarrow -0.5516389180 a^3 + 4.432882415 a^2$$
 $+ 0.000005306971905 a^2$
 $\sqrt{1.025000000 \cdot 10^{10} a^2 - 6.000000000 \cdot 10^{10} a + 8.780487804 \cdot 10^{10}}$
 $- 9.319560420 a$
 $- 0.00002497398544 a$
 $\sqrt{1.025000000 \cdot 10^{10} a^2 - 6.000000000 \cdot 10^{10} a + 8.780487804 \cdot 10^{10}}$
 $+ 1.652242329 \cdot 10^{-8}$
 $+ 1.290322580 \cdot 10^{-14}$
 $\sqrt{1.025000000 \cdot 10^{10} a^2 - 6.000000000 \cdot 10^{10} a + 8.780487804 \cdot 10^{10}}$

```
> s2:=solve(diff(S(a),a),a);
```

 $s2 := 1.306155359, 3.855134992, 11.56540468$

```
> a_max:=evalf(s2[1]);
```

 $a_{\max} := 1.306155359$

```
> evalf(x0);
```

 2.926829268

(9.15)

(9.16)

(9.17)

(9.18)